



粒子物理

7. 媒介粒子传播的相互作用和量子电动力学



曹庆宏

北京大学物理学院




Recap

★ Working towards a proper calculation of decay and scattering processes

Initially concentrate on:

- $e^+e^- \rightarrow \mu^+\mu^-$
- $e^-q \rightarrow e^-q$



▲ Lectures 3-4 covered the relativistic calculation of particle decay rates and cross sections

$$\sigma \propto \frac{|M|^2}{\text{flux}} \times (\text{phase space})$$

▲ Lecture 5 covered relativistic treatment of spin-half particles

Dirac Equation

▲ This lecture concentrates on the **Lorentz Invariant Matrix Element**

- Interaction by particle exchange
- Introduction to Feynman diagrams
- The Feynman rules for QED

Interaction by Particle Exchange

- Calculate transition rates from Fermi's Golden Rule

$$\Gamma_{fi} = 2\pi |T_{fi}|^2 \rho(E_f)$$

where T_{fi} is perturbation expansion for the Transition Matrix Element


$$T_{fi} = \langle f|V|i\rangle + \sum_{j \neq i} \frac{\langle f|V|j\rangle \langle j|V|i\rangle}{E_i - E_j} + \dots$$

- For particle scattering, the first two terms in the perturbation series can be viewed as:

“scattering in a potential”




“scattering via an intermediate state”



- “Classical picture” – particles act as sources for fields which give rise a potential in which other particles scatter – “action at a distance”
- “Quantum Field Theory picture” – forces arise due to the exchange of virtual particles. No action at a distance + **forces** between particles now **due to particles**

- Consider the particle interaction $a + b \rightarrow c + d$ which occurs via an intermediate state corresponding to the exchange of particle x
- One possible space-time picture of this process is:



Initial state i : $a + b$

Final state f : $c + d$

Intermediate state j : $c + b + x$

- This time-ordered diagram corresponds to a “emitting” x and then b absorbing x


- The corresponding term in the perturbation expansion is:

$$T_{fi} = \frac{\langle f|V|j\rangle\langle j|V|i\rangle}{E_i - E_j}$$

$$T_{fi}^{ab} = \frac{\langle d|V|x+b\rangle\langle c+x|V|a\rangle}{(E_a + E_b) - (E_c + E_x + E_b)}$$

- T_{fi}^{ab} refers to the time-ordering where a emits x before b absorbs it

- Need an expression for $\langle c+x|V|a\rangle$ in non-invariant matrix element T_{fi}



- Ultimately aiming to obtain Lorentz Invariant ME

- Recall T_{fi} is related to the invariant matrix element by

$$T_{fi} = \prod_k (2E_k)^{-1/2} M_{fi}$$

where k runs over all particles in the matrix element

- Here we have $\langle c+x|V|a\rangle = \frac{M_{(a \rightarrow c+x)}}{(2E_a 2E_c 2E_x)^{1/2}}$

$M_{(a \rightarrow c+x)}$ is the “**Lorentz Invariant**” matrix element for $a \rightarrow c+x$

- ★ The simplest Lorentz Invariant quantity is a scalar, in this case

$$\langle c+x|V|a\rangle = \frac{g_a}{(2E_a 2E_c 2E_x)^{1/2}}$$

g_a is a measure of the strength of the interaction $a \rightarrow c+x$

Note : the matrix element is only LI in the sense that it is defined in terms of LI wave-function normalisations and that the form of the coupling is LI


Note : in this “illustrative” example g is not dimensionless.

Similarly

$$\langle d|V|x+b\rangle = \frac{g_b}{(2E_b 2E_d 2E_x)^{1/2}}$$

Giving

$$\begin{aligned} T_{fi}^{ab} &= \frac{\langle d|V|x+b\rangle \langle c+x|V|a\rangle}{(E_a + E_b) - (E_c + E_x + E_b)} \\ &= \frac{1}{2E_x} \cdot \frac{1}{(2E_a 2E_b 2E_c 2E_d)^{1/2}} \cdot \frac{g_a g_b}{(E_a - E_c - E_x)} \end{aligned}$$




★ The “Lorentz Invariant” matrix element for the **entire** process is

$$\begin{aligned} M_{fi}^{ab} &= (2E_a 2E_b 2E_c 2E_d)^{1/2} T_{fi}^{ab} \\ &= \frac{1}{2E_x} \cdot \frac{g_a g_b}{(E_a - E_c - E_x)} \end{aligned}$$



Note:

- ◆ M_{fi}^{ab} refers to the time-ordering where a emits x before b absorbs it
It is **not Lorentz invariant**, order of events in time depends on frame
- ◆ Momentum is conserved at each interaction vertex but not energy
 $E_j \neq E_i$
- ◆ Particle x is “on-mass shell” i.e. $E_x^2 = \vec{p}_x^2 + m^2$

★ But need to consider also the other time ordering for the process



- This time-ordered diagram corresponds to ***b*** “emitting” \tilde{x} and then ***a*** absorbing \tilde{x}
- \tilde{x} is the anti-particle of x e.g.



• The Lorentz invariant matrix element for this time ordering is:

$$M_{fi}^{ba} = \frac{1}{2E_x} \cdot \frac{g_a g_b}{(E_b - E_d - E_x)}$$

★ Sum over matrix elements corresponding to same final state:

$$\begin{aligned} M_{fi} &= M_{fi}^{ab} + M_{fi}^{ba} \\ &= \frac{g_a g_b}{2E_x} \cdot \left(\frac{1}{E_a - E_c - E_x} + \frac{1}{E_b - E_d - E_x} \right) \\ &= \frac{g_a g_b}{2E_x} \cdot \left(\frac{1}{E_a - E_c - E_x} - \frac{1}{E_a - E_c + E_x} \right) \end{aligned}$$

Energy conservation:
($E_a + E_b = E_c + E_d$)


- Which gives
$$M_{fi} = \frac{g_a g_b}{2E_x} \cdot \frac{2E_x}{(E_a - E_c)^2 - E_x^2}$$

$$= \frac{g_a g_b}{(E_a - E_c)^2 - E_x^2}$$

- From 1st time ordering $E_x^2 = \vec{p}_x^2 + m_x^2 = (\vec{p}_a - \vec{p}_c)^2 + m_x^2$

giving
$$M_{fi} = \frac{g_a g_b}{(E_a - E_c)^2 - (\vec{p}_a - \vec{p}_c)^2 - m_x^2}$$

$$= \frac{g_a g_b}{(p_a - p_c)^2 - m_x^2}$$





→
$$M_{fi} = \frac{g_a g_b}{q^2 - m_x^2}$$

- After summing over all possible time orderings, M_{fi} is (as anticipated) **Lorentz invariant**. This is a remarkable result – the sum over all time orderings gives a frame independent matrix element.
- Exactly the same result would have been obtained by considering the annihilation process

Feynman Diagrams

- The sum over all possible time-orderings is represented by a **FEYNMAN diagram**




In a Feynman diagram:

- the LHS represents the initial state
- the RHS is the final state
- everything in between is “how the interaction happened”

- It is important to remember that **energy and momentum** are conserved at each interaction vertex in the diagram.
- The factor $1/(q^2 - m_x^2)$ is the propagator; it arises naturally from the above discussion of interaction by particle exchange

★ The matrix element: $M_{fi} = \frac{g_a g_b}{q^2 - m_x^2}$ depends on:

- The fundamental strength of the interaction at the two vertices g_a, g_b
- The four-momentum, q , carried by the (virtual) particle which is determined from energy/momentum conservation at the vertices.
Note q^2 can be either positive or negative.



Here $q = p_1 - p_3 = p_4 - p_2 = t$


“t-channel”

For **elastic scattering**: $p_1 = (E, \vec{p}_1)$; $p_3 = (E, \vec{p}_3)$

$$q^2 = (E - E)^2 - (\vec{p}_1 - \vec{p}_3)^2$$

$$q^2 < 0$$

termed “space-like”



Here $q = p_1 + p_2 = p_3 + p_4 = s$

“s-channel”

In CoM: $p_1 = (E, \vec{p})$; $p_2 = (E, -\vec{p})$


$$q^2 = (E + E)^2 - (\vec{p} - \vec{p})^2 = 4E^2$$

$$q^2 > 0$$


termed “time-like”

Virtual Particles

“Time-ordered QM”



Feynman diagram



$$M_{fi} = \frac{g_a g_b}{q^2 - m_x^2}$$

- Momentum conserved at vertices
- Energy **not** conserved at vertices
- Exchanged particle “**on mass shell**”

$$E_x^2 - |\vec{p}_x|^2 = m_x^2$$

- Momentum **AND** energy conserved at interaction vertices
- Exchanged particle “**off mass shell**”


VIRTUAL PARTICLE

- Can think of observable “on mass shell” particles as propagating waves and unobservable virtual particles as normal modes between the source particles:



Aside: $V(r)$ from Particle Exchange


- ★ Can view the scattering of an electron by a proton at rest in two ways:
 - Interaction by particle exchange in 2nd order perturbation theory.



$$M_{fi} = \frac{g_a g_b}{q^2 - m_x^2}$$

- Could also evaluate the same process in first order perturbation theory treating proton as a fixed source of a field which gives rise to a potential $V(r)$

$$M = \langle \psi_f | V(r) | \psi_i \rangle$$



Obtain same expression for M_{fi} using

$$V(r) = g_a g_b \frac{e^{-mr}}{r} \quad \text{YUKAWA potential}$$

- ★ In this way can relate potential and forces to the particle exchange picture
- ★ However, scattering from a fixed potential $V(r)$ is not a relativistic invariant view

Quantum Electrodynamics (QED)

- ★ Now consider the interaction of an electron and tau lepton by the exchange of a photon. Although the general ideas we applied previously still hold, we now have to account for the **spin of the electron/tau-lepton** and also the **spin (polarization) of the virtual photon**.

- The basic interaction between a photon and a charged particle can be introduced by making the **minimal substitution**

$$\vec{p} \rightarrow \vec{p} - q\vec{A}; \quad E \rightarrow E - q\phi \quad \text{(here } q = \text{charge)}$$

In QM: $\vec{p} = -i\vec{\nabla}; \quad E = i\partial/\partial t$

Therefore make substitution: $i\partial_\mu \rightarrow i\partial_\mu - qA_\mu$

where $A_\mu = (\phi, -\vec{A}); \quad \partial_\mu = (\partial/\partial t, +\vec{\nabla})$

- The Dirac equation:

$$\gamma^\mu \partial_\mu \psi + im\psi = 0 \quad \rightarrow \quad \gamma^\mu \partial_\mu \psi + iq\gamma^\mu A_\mu \psi + im\psi = 0$$

$$(\times i) \quad \rightarrow \quad i\gamma^0 \frac{\partial \psi}{\partial t} + i\vec{\gamma} \cdot \vec{\nabla} \psi - q\gamma^\mu A_\mu \psi - m\psi = 0$$

$$i\gamma^0 \frac{\partial \psi}{\partial t} = \gamma^0 \hat{H} \psi = m\psi - i\vec{\gamma} \cdot \vec{\nabla} \psi + q\gamma^\mu A_\mu \psi$$

$$\times \gamma^0 : \quad \hat{H} \psi = \underbrace{(\gamma^0 m - i\gamma^0 \vec{\gamma} \cdot \vec{\nabla})}_{\text{Combined rest mass + K.E.}} \psi + \underbrace{q\gamma^0 \gamma^\mu A_\mu}_{\text{Potential energy}} \psi$$

- We can identify the potential energy of a charged spin-half particle in an electromagnetic field as:

$$\hat{V}_D = q\gamma^0 \gamma^\mu A_\mu$$

(note the A_0 term is just: $q\gamma^0 \gamma^0 A_0 = q\phi$)

- The final complication is that we have to account for the photon polarization states.

$$A_\mu = \epsilon_\mu^{(\lambda)} e^{i(\vec{p} \cdot \vec{r} - Et)}$$

e.g. for a real photon propagating in the z direction we have two orthogonal transverse polarization states

$$\epsilon^{(1)} = \begin{pmatrix} 0 \\ 1 \\ 0 \\ 0 \end{pmatrix} \quad \epsilon^{(2)} = \begin{pmatrix} 0 \\ 0 \\ 1 \\ 0 \end{pmatrix}$$

Could equally have chosen circularly polarized states

- Previously with the example of a simple spin-less interaction we had:

$$M = \langle \psi_c | V | \psi_a \rangle \frac{1}{q^2 - m_x^2} \langle \psi_d | V | \psi_b \rangle$$

\parallel
 g_a
 \parallel
 g_b


- ★ In QED we could again go through the procedure of summing the time-orderings using Dirac spinors and the expression for \hat{V}_D . If we were to do this, remembering to sum over all photon polarizations, we would obtain:

$$M = [u_e^\dagger(p_3) q_e \gamma^0 \gamma^\mu u_e(p_1)] \sum_\lambda \frac{\epsilon_\mu^\lambda (\epsilon_\nu^\lambda)^*}{q^2} [u_\tau^\dagger(p_4) q_\tau \gamma^0 \gamma^\nu u_\tau(p_2)]$$

Interaction of e^- with photon

Massless photon propagator summing over polarizations

Interaction of τ^- with photon



- All the physics of **QED** is in the above expression !

- The sum over the polarizations of the **VIRTUAL** photon has to include longitudinal and scalar contributions, i.e. 4 polarization states

$$\epsilon^{(0)} = \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix} \quad \epsilon^{(1)} = \begin{pmatrix} 0 \\ 1 \\ 0 \\ 0 \end{pmatrix} \quad \epsilon^{(2)} = \begin{pmatrix} 0 \\ 0 \\ 1 \\ 0 \end{pmatrix} \quad \epsilon^{(3)} = \begin{pmatrix} 0 \\ 0 \\ 0 \\ 1 \end{pmatrix}$$

and gives:

$$\sum_{\lambda} \epsilon_{\mu}^{\lambda} (\epsilon_{\nu}^{\lambda})^* = -g_{\mu\nu}$$

This is not obvious – for the moment just take it on trust

and the invariant matrix element becomes:

$$M = [u_e^{\dagger}(p_3) q_e \gamma^0 \gamma^{\mu} u_e(p_1)] \frac{-g^{\mu\nu}}{q^2} [u_{\tau}^{\dagger}(p_4) q_{\tau} \gamma^0 \gamma^{\nu} u_{\tau}(p_2)]$$

- Using the definition of the adjoint spinor $\bar{\psi} = \psi^{\dagger} \gamma^0$

$$M = [\bar{u}_e(p_3) q_e \gamma^{\mu} u_e(p_1)] \frac{-g^{\mu\nu}}{q^2} [\bar{u}_{\tau}(p_4) q_{\tau} \gamma^{\nu} u_{\tau}(p_2)]$$

- ★ This is a remarkably simple expression ! It is shown in Appendix V of Handout 2 that $\bar{u}_1 \gamma^{\mu} u_2$ transforms as a four vector. Writing

$$j_e^{\mu} = \bar{u}_e(p_3) \gamma^{\mu} u_e(p_1) \quad j_{\tau}^{\nu} = \bar{u}_{\tau}(p_4) \gamma^{\nu} u_{\tau}(p_2)$$

$$M = -q_e q_{\tau} \frac{j_e \cdot j_{\tau}}{q^2} \quad \text{showing that } M \text{ is Lorentz Invariant}$$


Feynman Rules for QED

- It should be remembered that the expression

$$M = [\bar{u}_e(p_3)q_e\gamma^\mu u_e(p_1)] \frac{-g^{\mu\nu}}{q^2} [\bar{u}_\tau(p_4)q_\tau\gamma^\nu u_\tau(p_2)]$$

hides a lot of complexity. We have summed over all possible **time-orderings** and summed over all **polarization states** of the virtual photon. If we are then presented with a new Feynman diagram we don't want to go through the full calculation again. Fortunately this isn't necessary – can just write down matrix element using a set of simple rules

Basic Feynman Rules:



- Propagator factor for each internal line
(i.e. each internal virtual particle)
- Dirac Spinor for each external line
(i.e. each real incoming or outgoing particle)
- Vertex factor for each vertex

Basic Rules for QED

External Lines

spin 1/2		incoming particle	$u(p)$	
		outgoing particle	$\bar{u}(p)$	
		incoming antiparticle	$\bar{v}(p)$	
		outgoing antiparticle	$v(p)$	
spin 1		incoming photon	$\epsilon^\mu(p)$	
		outgoing photon	$\epsilon^\mu(p)^*$	

Internal Lines (propagators)

spin 1 photon

spin 1/2 fermion


$$-\frac{ig_{\mu\nu}}{q^2}$$


$$\frac{i(\gamma^\mu q_\mu + m)}{q^2 - m^2}$$


Vertex Factors


spin 1/2 fermion (charge $-|e|$)

$$ie\gamma^\mu$$



Matrix Element $-iM$ = product of all factors

e.g.



$$\bar{u}_e(p_3)[ie\gamma^\mu]u_e(p_1)$$


$$\frac{-ig_{\mu\nu}}{q^2}$$

$$\bar{u}_\tau(p_4)[ie\gamma^\nu]u_\tau(p_2)$$

$$-iM = [\bar{u}_e(p_3)ie\gamma^\mu u_e(p_1)] \frac{-ig_{\mu\nu}}{q^2} [\bar{u}_\tau(p_4)ie\gamma^\nu u_\tau(p_2)]$$

- Which is the same expression as we obtained previously

e.g.



$$-iM = [\bar{\nu}(p_2)ie\gamma^\mu u(p_1)] \frac{-ig_{\mu\nu}}{q^2} [\bar{u}(p_3)ie\gamma^\nu \nu(p_4)]$$

Note:

- ◆ At each vertex the adjoint spinor is written first
- ◆ Each vertex has a different index
- ◆ The $g_{\mu\nu}$ of the propagator connects the indices at the vertices

Summary

- ★ Interaction by particle exchange naturally gives rise to **Lorentz Invariant Matrix Element** of the form

$$M_{fi} = \frac{g_a g_b}{q^2 - m_x^2}$$

- ★ Derived the basic interaction in **QED** taking into account the spins of the fermions and polarization of the virtual photons:

$$-iM = [\bar{u}(p_3)ie\gamma^\mu u(p_1)] \frac{-ig_{\mu\nu}}{q^2} [\bar{u}(p_4)ie\gamma^\nu u(p_2)]$$

- ★ **We now have all the elements to perform proper calculations in QED !**